

THE CRUCIAL ROLE OF FORENSIC BOTANY IN THE SOLUTION OF JUDICIAL CASES

MARINA MORABITO ^{a*} AND ROBERTA SOMMA ^{bc}

ABSTRACT. A few years ago, two subjects disappeared and were found lifeless in the surrounding countryside. Understanding the causes of their death was particularly complex for the state of conservation of the human remains. A multi-disciplinary approach allowed experts in botany and geology to identify some algae traces on victims' belongings. Observations of microalgae, in association with other geological and botanical evidence, allowed investigators to geolocate the passage of one of the victims and to exclude that the victims' belongings were submerged under water. The present investigation proved how a traditional microscopic approach, based on the identification of associations of morpho-types of microalgae, could be decisive for linking the presence of one of the victims to specific sites of the territory where the body was found lifeless.

1. Introduction

The principle of Locard states that when two items or bodies interact, an inevitable contact responsible for a transfer of traces occurs. This principle, at the base of all forensic sciences, allows investigators to link a suspect to a victim or to a crime scene, because of the transfer of physical evidence. In outdoor crime scenes, in the countryside, geological and soil evidence may be represented by inorganic and organic traces. These latter may be animal and vegetal. Vegetal traces present in the soil evidence may be studied by forensic botanists in team with geologists, providing very useful information as demonstrated by Lombardi (1999) for the case of the kidnapping and homicide of Aldo Moro. Forensic botanists may investigate forensic traces of vascular and non-vascular plants. Among the non-vascular plants (*i.e.*, they do not have roots, stems or leaves), algae are unicellular or multicellular organisms widely distributed in all aquatic environments or simply in humid places, such as trees, walls, puddles, or moist soils (Sheath and Wehr 2015; Graham *et al.* 2019). Therefore, it is very likely that objects or people interacting in or near aquatic or wet environments encounter algae, which consequently become organisms potentially useful for linking a subject or an object to a site (D. W. Hall 2012; Hard and Wallace 2015; Levin *et al.* 2017; Magni *et al.* 2020). Aquatic ecologists have long recognized the critical importance of algae to the functioning of these ecosystems, as well as the role they play as determinants or indicators of water quality within them (Stevenson *et al.* 1996). It is this strong ecological

foundation that led scientists to explore their role in forensic investigations (Keiper and Casamatta 2001; Cameron 2004; Scott *et al.* 2014, 2017; Thakar, Luthra, and Khattar 2018; Wanner *et al.* 2018; Spikmans 2019; Scott *et al.* 2021; M. K. Sharma, A. Sharma, and Bhatt 2021; I. Bogusz, M. Bogusz, and Elazna-Wieczorek 2022). Due to the prominent role and prevalence of algae in aquatic ecosystems, it is very likely that objects or people involved in an accident or crime that occurs in water (even partially as in a puddle) may preserve traces of algae transferred from the surrounding environments. Peculiar species of algae exist in certain terrestrial environments such as shaded tree bark (*e.g.*, the north side in the northern hemisphere), shaded house facades or in the soil (Fitzpatrick and Donnelly 2021). Due to their prevalence and importance in nature, algae have a high probability of being a significant element in a forensic investigation. In cases of deaths in freshwaters, or in their proximity, the similarity of specific algae found as traces on clothing, vehicle or persons with the algal species growing at the scene of investigated events, or where physical evidence (such as a weapon), can provide valuable circumstantial evidence of correlation (Coyle *et al.* 2005). The comparison is possible between distinctive combinations of one or two species of indicator algae or, alternatively, between algal assemblages (*i.e.*, groups of species) that may not be typical of a defined location, but rather to a particular type of water body (*e.g.*, freshwater rather than marine, streams rather than ponds, or waters near sewers or sources of thermal pollution, *etc.*) (Britton and Greeson 1989; Siver, Lord, and McCarthy 1994; Hard and Wallace 2015). Furthermore, the presence or lack of algae in body organs (such as lungs) can be used for positive or negative diagnosis of drowning death, and the relative abundances of diatoms (*i.e.* *microalgae* with a cell wall composed of silica) in such tissues can be used to infer the probable location of the drowning (Pollanen, Cheung, and Chiasson 1997; Pollanen 1998; Zimmerman and Wallace 2008; Diaz-Palma *et al.* 2009; Saini, Khanagwal, and Singh 2017; Liu *et al.* 2020; Zhou *et al.* 2020; Du *et al.* 2022). Differently for land plants, whose anatomy is rich in providing diagnostic characteristics (Ladd and H. C. Lee 2005; Aquila *et al.* 2018; Alotaibi *et al.* 2020), both vegetative and reproductive, algae in general, and especially microalgae, are tricky to identify at the species level (R. E. Lee 2018). Classic taxonomic identifications have focused on morphological characters observed at the light microscopy (Palmer 1959; Bellinger and Sigeo 2015). However, modern research showed wide phenotypic plasticity and cryptic diversity and pointed on a rather different phylogenetic clustering (Verbruggen and Theriot 2008). A modern taxonomic approach generally does not fail of including ultrastructural (Scanning Electron Microscope - SEM and Transmission Electron Microscopy - TEM), phylogenetic, and phylogenomics data (Bhattacharya and Price 2020). However, in several professional contexts, such methods are not applicable and morphological identification at the light microscope is the standard for field samples (Cox and Cox 1996; Dillard 1999; Wehr, Sheath, and Kociolek 2015). The aim of the present paper was to verify whether a traditional morphological approach based on taxonomic identification of algae could be useful for exclusionary/comparative forensic purposes, bringing information for the comparison between samples of unknown origin with samples of known origin (Somma 2023c) related to a casework. An *atlas* of morphotypes (tentative *taxa*) recorded for each sample (both of known and unknown origin) was produced. A comparison was made of morphotypes identified in the samples of unknown origin to those of known origin and the peculiar associations of taxonomic entities in each sample

was used to evaluate similarity among traces found on the victim' shoes and each collection site.

2. Forensic Botany

The methodologies and principles of botany can be applied in the investigation of criminal cases being one of the forensic sciences applied to criminal law (Picozzi and Intini 2009; D. W. Hall 2012; Saferstein 2017). Both objects or persons present in the scene of events can be linked to the scene itself thanks to the contact, and the consequent gained traces, with fragments, individuals or populations of land plants, algae or fungi (Coyle *et al.* 2005; Hard and Wallace 2015). Botanical elements useful in the forensic investigations are seeds, fruits, stems, leaves, roots, algal thalli, or fragments of any of them, both macro and microscopic, as well as in the latter dimensional scale, pollens, spores, and microalgae, which are mixed in the soil as its organic component (Ladd and H. C. Lee 2005) in continental environments. To either connect or dissociate victims to suspects or crimes scenes, the transfer of plant elements from the environment surrounding the victim or the suspect, especially from the ground, can be useful on any occasion, but inevitably in events that took place outdoors (Ladd and H. C. Lee 2005). In circumstances where the scene of events is known, it is advisable to produce a reference collection to be used for comparison with samples of unknown origin found on objects or persons (Ladd and H. C. Lee 2005). The questioned samples (unknown samples), when it is possible, are the first specimens to be examined for the taxonomic identification of the plant species present. After this phase, the same examination is usually carried out on the samples collected from the sites of the events of investigative interest (known samples). In addition, forensic botany also relies on anatomical, morphological, palynological, ecological, and vegetational observations (*e.g.*, Horswell *et al.* 2002; Coyle *et al.* 2005; D. W. Hall 2012). More commonly, botanical traces provide potential class evidence, but occasionally, for example thanks to molecular analyses (DNA analyses), individual evidence can be obtained (*sensu* Saferstein 2017).

3. Case Study

A few years ago, two subjects (victims 1 and 2) disappeared and were found lifeless in the countryside. Understanding the causes of their death was particularly complex for the state of conservation of the human remains. The investigating authority tried not to neglect any element useful for the reconstruction of the events and consequently technical (Somma 2023b) investigations on the inorganic and organic traces found on the victims were also ordered (Somma *et al.* 2023c). Because events occurred outdoor, many vegetal traces were present on the victims' bodies and their belongings. These traces, showing potentially useful organisms for forensic investigation, were carefully analysed.

3.1. Materials and Methods. Geological and botanical traces collected from the clothing and footwear finds of the victims (samples of unknown origin) were scanned both macroscopically and at the stereomicroscope to check the presence of possible algae. Aggregates of microalgae were also sampled from moist soil and fresh water from ponds collected on the sites under investigation (samples of known origin) (Figure 1) and fixed in formalin at the sampling site (final concentration at 4%). All samples were observed for comparative

purposes at the light microscopy (Laborlux 12, 40-1000 [o.i.] magnifications, LEITZ, Germany, equipped with a digital camera 12 MP, Apple Inc., USA). Microalgae were separated from the soil by dispersing them in sterile Petri dishes with distilled water and then small aliquots were mounted on microscope slides. Taxonomic identification of the microalgae was performed on a morphological basis¹ (Prescott, Croasdale, and Vinyard 1964; John *et al.* 2002; Bellinger and Sigeo 2015; Wehr, Sheath, and Kociolek 2015). Where identification was not achievable at the species level, the smallest identifiable taxonomic level was recorded. Morphological characteristics (shape, size, color, taking into consideration the different state of conservation of the finds) and cellular characteristics (wall, unicellular, colonial, multicellular organization) were carefully observed. SEM images (microscope SEM QUANTA FEG 450, FEI Europe, Netherlands) were also taken and observed to scan for the presence of microalgae in the clothing of the victims.

4. Results

An aggregate mass mixed to geological evidence traces, a few millimeters in size, was found by the forensic geologist (R.S.) stuck in the sole of the shoe of victim 1 (unknown sample) (Figure 1A). It was characterized by a greenish color, but its plastic behaviour suggested that the material was not inorganic. This aggregate, observed by the botanist (M.M.), revealed to be made of a microalgal assemblage (Figure 1B). No algae were recorded on other part of the shoes or on victim 1's clothes.




FIGURE 1. A) Close up of the sole of the shoe of victim 1 with the microalgal aggregate before sampling (upper) and after (lower). B) Algae isolated from the aggregate (unknown sample) at the light microscope.

¹M. D. Guiry and G. M. Guiry, "AlgaeBase. World-Wide Electronic Publication" National University of Ireland, Galway, 2023. URL: <http://www.algaebase.org>

Five distinct species of green algae (*Chlorophyta*), one morphotype of blue algae (*Cyanophyta*), determined at the level of order, and two diatoms (*Ochrophyta*), determined at the level of order, for a total of seven different taxonomic entities were identified on the unknown sample collected on the shoe of victim 1 (Table 1).

TABLE 1. Algae identified in the unknown sample of soil isolated from the sole of the shoe of victim 1.

Taxonomy

Deasonnia sp. (*Chlamydomonadales*, *Chlorophyta*)

Deasonnia sp. young cells, or other (*Chlamydomonadales*, or *Chlorococcales* (*Chlorophyta*))

Apatococcus sp. (*Chlorellales*, *Chlorophyta*)

Apatococcus sp. (*Chlorellales*, *Chlorophyta*)

Species of *Chlorellales*, or *Chlamydomonadales* (*Chlorophyta*)

Scenedesmus sp. (*Sphaeropleales*, *Chlorophyta*)

Desmodesmus sp. (*Sphaeropleales*, *Chlorophyta*)

Chroococcales (*Cyanophyta*) in complex with *Chlorophyta*

Diatoms (*Pennales*, *Ochrophyta*) (a more detailed identification is not possible as cells were in girdle view)

A single diatom (*Pennales*, *Ochrophyta*), determined at the level of order, was found in a SEM graph of the sole of a shoe of victim 2 (unknown sample) (Figure 2). No other types of algae were recorded on other part of the shoe or on victim 2's clothes.




FIGURE 2. Diatom isolated from the sole of the shoe of victim 2 at the SEM (unknown sample).

Eight distinct species of green algae (five *Chlorophyta* and three *Streptophyta*), two morphotypes of blue algae (*Cyanophyta*), determined at the level of order, and several

morphotypes of diatoms (*Ochrophyta*), determined at the level of order, for a total of eleven different taxonomic entities were identified in distinct associations in samples of known origin collected on the scene of events in freshwaters and ponds (Table 2; Figure 3).

TABLE 2. Algae identified in samples of moist soil and water from ponds collected on the investigated area (samples of known origin).

Taxonomy

Deasonnia sp. (*Chlamydomadales*, *Chlorophyta*)

Deasonnia sp. young cells, or other (*Chlamydomadales*, or *Chlorococcales* (*Chlorophyta*))

Apatococcus sp. (*Chlorellales*, *Chlorophyta*)

Chroococcales (*Cyanophyta*) in complex with (*Chlorophyta*)

Clasterium sp. 1 (*Desmidiiales*, *Streptophyta*)

Clasterium sp. 2 (*Sphaeropleales*, *Chlorophyta*)

Diatoms (*Pennales*, *Ochrophyta*)

Filamentous *Cyanophyta*

Scenedesmus sp. (*Sphaeropleales*, *Chlorophyta*)

Species of *Chlorellales* o *Chlamydomadales* (*Chlorophyta*)

Spirogyra sp. (*Zygnematales*, *Streptophyta*)




FIGURE 3. A) The countryside surrounding the site of the discovery of the victims' bodies with the indication of the puddle of Figure 1 B (star); B) the mud of the puddle (known sample) where algae thrive, similar to those found in the shoes of victim 1 (unknown sample).

4.1. Description of taxa identified in the unknown and known samples. *Deasonia* sp. (*Chlamydomadales*, *Chlorophyta*), *Deasonia* sp. young cells, or other *Chlamydomadales* or *Chlorococcales* (*Chlorophyta*) (Figure 4 A-B).

Species of the genus *Deasonia* Ettl et Komárek have single cells (plurinucleate); young cells are elliptical, ovoid, while mature cells are always spherical with a thickened cell wall. They have parietal chloroplasts with bifurcations and indentations in mature cells, finally

reticulated, while cup-shaped perforated in young cells; in young cells pyrenoid is lateral in compact chloroplasts and moves to the center of the chloroplast in mature cells; vegetative stages are multinucleated, they can form biflagellate zoospores; the species are present in moist soil (Shubert and Gärtner 2015). The young cells have a more generic morphology which can be confused with other species of the same order (*Chlamydomonadales*) or of the order *Chlorococcales* (*Chlorophyta*).

Apatococcus sp. (*Chlorellales*, *Chlorophyta*) (Figure 4 C)

Species of the genus *Apatococcus* Brand *em.* Geitler have cells rounded to spherical or slightly flattened at the corners, more or less irregular cell packets formed by divisions in two or three consecutive perpendicular directions or multilayered pseudoparenchymatous cell aggregates, occasionally the beginning of the formation of short filaments can be observed; cell wall with smooth to rough surface, somewhat thickened in older cells; chloroplast in young parietal cells, in older cells partially constricted in the center, lobed, lacking pyrenoid (Shubert and Gärtner 2015). Rarely reported, but probably very common or cosmopolitan, the actual distribution is poorly known (Shubert and Gärtner 2015).

Species of *Chlorellales* or *Chlamydomonadales* (*Chlorophyta*) (Figure 4 D)

There are over 200 recognized genera among the non-motile and colonial coccoid green algae, with a range of morphologies and ecological habitats (Shubert and Gärtner 2015).

Scenedesmus sp. (*Sphaeropleales*, *Chlorophyta*) (Figure 4 E)

Species of the genus *Scenedesmus* Meyen form colonies of 2, 4, 8 or 16 (rarely 32) cells, flattened, with long axes of parallel cells, laterally adjacent, arranged in a single linear or alternating series, ellipsoidal, ovoid or crescent or tapering towards each end, cell wall spineless, parietal chloroplast, usually with a pyrenoid (Shubert and Gärtner 2015). Reported worldwide and frequently abundant in nutrient-rich (especially high inorganic N) waters; it is commonly found in association with other coccoid algae genera (spiny-producing species, such as those of the genus *Desmodesmus*) (Shubert and Gärtner 2015).

Desmodesmus sp. (*Sphaeropleales*, *Chlorophyta*) (Figure 4 F)

Species of the genus *Desmodesmus* An, Friedl et Hegewald form colonies of 2, 4, 8 or 16 cells or unicellular forms, with long axes of parallel cells, laterally flanked, arranged in a single linear or alternating series; cells ellipsoidal, to ovoid, spines usually present on terminal cells and/or medial cells, occasionally absent, parietal chloroplasts, usually with a pyrenoid (Shubert and Gärtner 2015). An extremely common (and occasionally abundant) genus in continental water bodies (Shubert and Gärtner 2015). *Desmodesmus* species that typically produce colonial forms may occasionally produce single cells depending on environmental conditions, such as phosphorus concentration (Trainor 2014).

Closterium sp. (*Desmidiiales*, *Streptophyta*) (Figure 4 G-H)

Species of the genus *Closterium* Nitzsch ex Ralfs have solitary cells, usually straight, arcuate, or lunate, many times longer than broad, in most species, tapering toward the apices, which may be pointed, broadly rounded, or truncated (J. D. Hall and McCourt 2015). Cells are unconstricted and may be inflated in the mid region in some species. Size varies by species from tens of micrometers in length to nearly a millimeter. In certain species, two semicells are visible due to intercalating wall material, with a central nucleus and two chloroplasts, axile and lobed with several pyrenoids. Sexual reproduction is known in several species, and zygospores may be irregularly globose or quadrate and are generally unornamented. Most species are benthic, loosely attached to sediments or on aquatic plants, but some are

planktonic. Some species of *Closterium* can be found in abundance on damp soils (Prescott *et al.* 1975).

Spirogyra sp. (*Zygnematales*, *Streptophyta*) (Figure 4 I)

Species of *Spirogyra* Link form long uniseriate filaments with cylindrical, unornamented, and unstricted cells, usually attached to substrata by rhizoids, resting on sediments or floating on the surface (J. D. Hall and McCourt 2015). Cells have one to several parietal, spiraling chloroplasts per cell, a central nucleus and a large central vacuole. Species reproduce both sexually via zygospores and asexually by vegetative reproduction and aplanospores. All spores have multilayered walls that may be variously colored and ornamented. Zygospores are useful for species-level identification but are rarely encountered in the field. The genus *Spirogyra* is one of the most widespread and common of all freshwater algae. Species are tolerant of a wide variety of environmental and chemical conditions, including some that are remarkably tolerant of pollutants and a few that tolerate brackish conditions.

Filamentous Cyanophyta (Figure 4 K-M)

Cyanophyta, or *Cyanobacteria*, occur in a wide range of morphologies and ecological forms, which colonize almost all habitats and are ubiquitous in aquatic and terrestrial communities, including almost all types of moist soils (Komárek and Johansen 2015a,b). In recent decades, ecological features, ultrastructural features, and molecular evidence have substantially influenced the knowledge and understanding of this group. As a result, the classification of species is undergoing radical changes. The cellular morphological approach to classification, regardless its considerable limits, is still used mainly for the identification of natural material (Komárek and Johansen 2015a). Most soil cyanobacteria are filamentous species, but several coccoid forms are also common. Filamentous cyanobacteria comprise some of the most widely recognized and important freshwater algae in the world, and their classification is equally complex and intersecting with that of the orders of coccoid representatives (Komárek and Johansen 2015b). Many of the filamentous *taxa* are important members of lake and riverine phytoplankton, while others form thick and dense beds in a wide variety of benthic habitats (Komárek and Johansen 2015b). Others form conspicuous masses floating on the water surface of various water bodies, or grow attached to stones, or epiphytes on aquatic angiosperms and other algae, epipellic (*i.e.*, on muddy sediments) and epipsammic (*i.e.*, on sand), and frequently colonize wet or damp lands (Komárek and Johansen 2015b).

Chroococcales (*Cyanophyta*) in complex aggregates with *Chlorophyta* (Figure 4 L)

Unicellular and colonial coccoid cyanobacteria were classified according to the traditional morphological system in the order *Chroococcales* (Komárek and Johansen 2015a). From modern investigations, mainly molecular and ultrastructural, it follows that coccoid cyanobacteria are a very heterogeneous group, developed in several distinct evolutionary lineages that led to a recent reclassification into phylogenetically distinct orders. Mucilaginous colonies of cyanobacteria can be spherical, oval, planar, or irregular (Komárek and Johansen 2015b). The structure of the sheaths and envelopes around the cells is different: mucilage can form a hyaline, amorphous mass, diffuse and marginal envelopes or structured and stratified sheaths (Komárek and Johansen 2015b). The gelatinous shells and sheaths (solid or structured outer layers) appear to perform a protective function in environments exposed to severe desiccation or solar radiation.




FIGURE 4. A) *Deasonia* sp. (*Chlamydomonadales*, *Chlorophyta*); B) *Deasonia* sp. young cells, or other *Chlamydomonadales* or *Chlorococcales* (*Chlorophyta*); C) *Apatococcus* sp. (*Chlorellales*, *Chlorophyta*); D) *Chlorellales* or *Chlamydomonadales* (*Chlorophyta*); E) *Scenedesmus* sp. (*Sphaeropleales*, *Chlorophyta*); F) *Desmodesmus* sp. (*Sphaeropleales*, *Chlorophyta*); G) *Closterium* sp. 1 (*Desmidiiales*, *Streptophyta*); H) *Closterium* sp. 2 (*Desmidiiales*, *Streptophyta*); I) *Spirogyra* sp. (*Zygnematales*, *Streptophyta*); J) diatoms (*Pennales*, *Ochrophyta*) in valve view; K, M) filamentous *Cyanophyta*; L) *Chroococcales* (*Cyanophyta*) in complex with *Chlorophyta*.

Diatoms (Pennales, Ochrophyta) (Figure 4 J)

Diatoms are unicellular algae present in almost all bodies of water, with a rigid cell wall composed of silica (SiO_2), known as a frustule, composed of two valves, a smaller hypovalve, which fits inside a larger epivalve, like two halves of a “box” (Kociolek, Spaulding, and Lowe 2015; Kociolek *et al.* 2015). Average cell size decreases in a population, because each valve produces a smaller complementary valve during cell division until it reaches a minimum size (usually one-third of maximum size) that induces sexual reproduction and restoration of maximum size (Edlund and Stoermer 1997). The siliceous walls of diatoms are often highly ornamented, and a large part of the taxonomy has always been based on their morphological characteristics, traditionally observed at the optical microscopy; currently, ultrastructural observations (SEM) of the frustule are considered essential for species identification, due to a strong morphological convergence between taxonomically distinct groups (Kociolek, Spaulding, and Lowe 2015). Among the diatoms, species with radial symmetry (*Centrales*) and others with bilateral symmetry (*Pennales*) are traditionally distinguished; among the latter, a subgroup has evolved a special structure involved in the movement of cells on surfaces, called raphe (Kociolek, Spaulding, and Lowe 2015). Modern phylogenetic systematics reorganizes these groups into several distinct lines. For simplicity, the *Centrales* and *Pennales* groupings had been used in this work, although we are aware that they are no longer taxonomically rigorous.

4.2. Comparative analyses. From the comparative analyses, a technical opinion of generic compatibility of the trace from the sole of the shoe of victim 1 (Table 1) with macroenvironments similar to the different freshwater points found in the area under investigation (Table 2) was expressed; in some site the peculiar association of microalgae were all present, in other ones only partially. Furthermore, a technical opinion of strong compatibility of the same sample of unknown origin with macro-environments similar to the bottom of a small artificial basin (known as “*gebbia*”) and to an ephemeral pool in the ground, respectively fed by a pipeline drawing spring water and its loss, was expressed. It was not possible to express a technical opinion of compatibility of the diatom observed in the micrograph of the sole of the shoe of victim 2 with any diatom observed in macroenvironments in the investigated area. Given the extremely small size of the observed diatom, it was not possible to exclude that the failure to find it might be due to difficulties to identify minute organisms in complex matrices, but also might depend on the insufficient observation time enforced by the investigations.

5. Discussion and Conclusions

Forensic sciences imply the use of expertise, principles, scientific methods, and modern technologies related to a wide range of disciplines, in order to fully investigate crimes and physical evidence that might be presented in a court of law (Picozzi and Intini 2009; Saferstein 2017). By definition, a “holistic” multidisciplinary approach is crucial for not undervaluing data and information from all traces both from the scenes of events and the persons involved (Somma 2022, 2023a). Especially in crimes occurring outdoor, collected evidence and analyses involving an array of disciplines, such as botany (Caccianiga *et al.* 2021; Morabito, Mondello, and Somma 2023; Somma *et al.* 2023c), geology (Ruffell and McKinley 2009; Somma *et al.* 2023c,d; Spoto, Barone, and Somma 2023), or paleontology (Di Maggio *et al.* 2014; Somma and Maniscalco 2023), are crucial in investigation where soils and plants are widespread. In addition, forensic botanists with geologists, geographers, coroners, and environmental psychologists may also collaborate in multidisciplinary teams in cases regarding the geographic profiling (Canter and Youngs 2003; Canter 2005; Somma 2023b) or searches for clandestine graves (Somma and Costa 2022; Byrd and Sutton 2023; Marra 2023; Somma and Costa 2023; Somma, Sutton, and Byrd 2023; Tagliabue *et al.* 2023), where also the plants may play a significant rule in dating the age of the burial. Other illicit activities involving plants may concern fake fossils of plants, frauds, and detection of counterfeit goods (Marra, Di Silvestro, and Somma 2023; Spoto 2023). In this paper, such multidisciplinary approach allowed the botany and geology experts to ascertain the presence of the two subjects in the scene of event and to reconstruct the path followed by them in their last hours of their lives. In particular, observations of microalgae, in association with all other geological and botanical evidence (Morabito, Mondello, and Somma 2023; Somma *et al.* 2023c), allowed investigators to geolocate the passage of victim 1 in a specific puddle located close to the disappearance site. Nowadays, taxonomic identification of microalgae often uses phylogenetic and ultrastructural approaches, which needs the involvement of skilled experts and complex techniques. Consequently, the before mentioned approach in forensic investigations is neglected, with few notable exceptions (*i.e.*, diatoms in drowning deaths, Magrey and Raj 2014; Zhao *et al.* 2016; Shen *et al.* 2019;

Zhang *et al.* 2020; Khurshid *et al.* 2021). Conversely, the present investigation proved how a traditional light microscopic approach could be decisive to associate known samples with unknown traces of soil, basing on the identification of associations of morpho-types of microalgae. Furthermore, a multidisciplinary approach, as it can be accomplished by a diversified working team, should be desirable in forensic investigations and achievable only when the education system includes forensic competencies in the *curricula* of studies (Spoto, Somma, and Crea 2021; Somma 2022; Baldino *et al.* 2023; Somma *et al.* 2023a,b). The large amount of data that can be collected by means of a holistic approach may allow to test carefully alternative hypotheses and, when concordant, contributes to reconstruct past events and dynamics. In complex cases, such as the casework here presented, a such approach may assist judicial authority also in the understanding of the reasons of certain criminal behaviours.

References

- Alotaibi, S. S., Sayed, S. M., Alosaimi, M., Alharthi, R., Banjar, A., Abdulqader, N., and Alhamed, R. (2020). "Pollen molecular biology: Applications in the forensic palynology and future prospects: A review". *Saudi journal of biological sciences* **27**(5), 1185–1190. DOI: [10.1016/j.sjbs.2020.02.019](https://doi.org/10.1016/j.sjbs.2020.02.019).
- Aquila, I., Gratteri, S., Sacco, M. A., and Ricci, P. (2018). "The role of forensic botany in solving a case: scientific evidence on the falsification of a crime scene". *Journal of forensic sciences* **63**(3), 961–964. DOI: [10.1111/1556-4029.13639](https://doi.org/10.1111/1556-4029.13639).
- Baldino, G., Ventura Spagnolo, E., Fodale, V., Pennisi, C., Mondello, C., Altadonna, A., Raffaele, M., Salmeri, F., Somma, R., Asmundo, A., and Sapienza, D. (2023). "The application of 3D virtual models in the judicial inspection of indoor and outdoor crime scenes". *Atti della Accademia Peloritana dei Pericolanti. Classe di Scienze Fisiche, Matematiche e Naturali* **101**(S1), A17. DOI: [10.1478/AAPP.101S1A17](https://doi.org/10.1478/AAPP.101S1A17).
- Bellingier, E. G. and Sigeo, D. C. (2015). *Freshwater algae: identification, enumeration and use as bioindicators*. John Wiley & Sons.
- Bhattacharya, D. and Price, D. C. (2020). "The algal tree of life from a genomics perspective". *Photosynthesis in algae: biochemical and physiological mechanisms*, 11–24.
- Bogusz, I., Bogusz, M., and Elazna-Wieczorek, J. (2022). "Diatoms from inland aquatic and soil habitats as indestructible and non removable forensic environmental evidence". *Journal of Forensic Sciences* **67**(4), 1490–1504. DOI: [10.1111/1556-4029.15017](https://doi.org/10.1111/1556-4029.15017).
- Britton, L. J. and Greeson, P. E. (1989). *Methods for collection and analysis of aquatic biological and microbiological samples*. US Government Printing Office. DOI: [10.3133/ofr88190](https://doi.org/10.3133/ofr88190).
- Byrd, J. H. and Sutton, L. (2023). "The Use of Forensic Entomology within Clandestine Gravesite Investigations". *Atti della Accademia Peloritana dei Pericolanti. Classe di Scienze Fisiche, Matematiche e Naturali* **101**(S1), A7. DOI: [10.1478/AAPP.101S1A7](https://doi.org/10.1478/AAPP.101S1A7).
- Caccianiga, M., Caccia, G., Mazzarelli, D., Salsarola, D., Poppa, P., Gaudio, D., Cappella, A., Franceschetti, L., Tambuzzi, S., Maggioni, L., and al., et (2021). "Common and much less common scenarios in which botany is crucial for forensic pathologist and anthropologists: a series of eight case studies". *International Journal of Legal Medicine* **135**, 1067–1077. DOI: [10.1007/s00414-020-02456-0](https://doi.org/10.1007/s00414-020-02456-0).
- Cameron, N. G. (2004). "The use of diatom analysis in forensic geoscience". *Geological Society, London, Special Publications* **232**(1), 277–280. DOI: [10.1144/GSL.SP.2004.232.01.25](https://doi.org/10.1144/GSL.SP.2004.232.01.25).
- Canter, D. V. (2005). *Mapping murder: Walking in killers' footsteps*. Virgin Books Publishers.
- Canter, D. V. and Youngs, D. (2003). "Forensic geology". In: *Handbook of psychology in legal contexts*. Ed. by R. C. Bull and C. D. V. Chichester, UK: Wiley, pp. 171–206.

- Cox, E. J. and Cox, E. J. (1996). *Identification of freshwater diatoms from live material*. Vol. 158. Chapman & Hall London. DOI: [10.2216/i0031-8884-36-4-330.1](https://doi.org/10.2216/i0031-8884-36-4-330.1).
- Coyle, H. M., Lee, C., Lin, W., Lee, H. C., and Palmbach, T. M. (2005). "Forensic botany: using plant evidence to aid in forensic death investigation". *Croatian medical journal* **46**(4), 606.
- Di Maggio, R. M., Barone, P. M., Pettinelli, E., Mattei, E., Lauro, S. E., and Banchelli, A. (2014). *Geologia Forense: Introduzione alle geoscienze applicate alle indagini giudiziarie*. Dario Flaccovio Editore.
- Díaz-Palma, P. A., Alucema, A., Hayashida, G., and Maidana, N. I. (2009). "Development and standardization of a microalgae test for determining deaths by drowning". *Forensic science international* **184**(1-3), 37–41. DOI: [10.1016/j.forsciint.2008.11.015](https://doi.org/10.1016/j.forsciint.2008.11.015).
- Dillard, G. E. (1999). *Common freshwater algae of the United States: An illustrated key to the genera (excluding the diatoms)*. Verlag. DOI: [97-8-344-350033-7](https://doi.org/10.1016/j.forsciint.2008.11.015).
- Du, Y., Liu, J., Li, Q., Li, H., Kang, X., Zheng, D., Shi, H., Xu, Q., Liu, C., Wang, H., and al., et (2022). "Concordance analysis of diatom types and patterns in lung tissue and drowning medium in laboratory animal model". *International journal of legal medicine* **136**(3), 911–917. DOI: [10.1007/s00414-021-02768-9](https://doi.org/10.1007/s00414-021-02768-9).
- Edlund, M. B. and Stoermer, E. F. (1997). "Ecological, Evolutionary, and Systematic Significance of Diatom Life Histories 1". *Journal of Phycology* **33**, 897–918. DOI: [10.1111/j.0022-3646.1997.00897.x](https://doi.org/10.1111/j.0022-3646.1997.00897.x).
- Fitzpatrick, R. W. and Donnelly, L. J. (2021). *An introduction to forensic soil science and forensic geology: a synthesis*. DOI: [10.1144/SP492-2021-8](https://doi.org/10.1144/SP492-2021-8).
- Graham, L. E., Graham, J. M., Wilcox, L. W., and Cook, M. E. (2019). *Algae (3rd Edition)*. LJLM Press.
- Hall, D. W. (2012). "Introduction to forensic botany". In: *Forensic botany: a practical guide*. Ed. by D. W. Hall and J. H. Byrd. John Wiley & Sons.
- Hall, J. D. and McCourt, R. M. (2015). "Conjugating green algae including desmids". In: *Freshwater Algae of North America*. Ed. by J. D. Wehr, R. G. Sheath, and J. P. Kociolek. Elsevier Inc. DOI: [10.1016/B978-0-12-385876-4.00009-8](https://doi.org/10.1016/B978-0-12-385876-4.00009-8).
- Hard, C. R. and Wallace, J. R. (2015). "Algae in forensic investigations". In: *Forensic botany: a practical guide*. Ed. by D. W. Hall and J. H. Byrd. John Wiley & Sons.
- Horswell, J., Cordiner, S., Maas, E., Martin, T., Sutherland, K., Speir, T., Nogales, B., and Osborn, A. (2002). "Forensic Comparison of Soils by Bacterial Community DNA Profiling". *Journal of forensic sciences* **47**, 350–3. DOI: [10.1520/JFS15256J](https://doi.org/10.1520/JFS15256J).
- John, D. M., Whitton, B. A., Brook, A. J., York, P. V., and Johnson, L. R. (2002). *The freshwater algal flora of the British Isles: An identification guide to freshwater and terrestrial algae*. Cambridge University Press.
- Keiper, J. B. and Casamatta, D. A. (2001). "Benthic organisms as forensic indicators". *Journal of the North American Benthological Society* **20**(2), 311–324. DOI: [10.2307/1468325](https://doi.org/10.2307/1468325).
- Khurshid, A., Shah, M. U., Khurshid, M., Sohail, A., Ali, G., Ali Sr, G., and al., et (2021). "Diatom-positive cadaver: drowning or homicide?" *Cureus* **13**(9), e18312. DOI: [10.7759/cureus.18312](https://doi.org/10.7759/cureus.18312).
- Kociolek, J. P., Spaulding, S. A., and Lowe, R. L. (2015). "Bacillariophyceae: The raphid diatoms". In: *Freshwater algae of North America*. Ed. by J. D. Wehr, R. G. Sheath, and J. P. Kociolek. Elsevier Inc, pp. 709–772. DOI: [10.1016/B978-0-12-385876-4.00016-5](https://doi.org/10.1016/B978-0-12-385876-4.00016-5).
- Kociolek, J. P., Theriot, E. C., Williams, D. M., Julius, M., Stoermer, E. F., and Kingston, J. C. (2015). "Centric and Araphid Diatoms". In: *Freshwater algae of North America*. Ed. by J. D. Wehr, R. G. Sheath, and J. P. Kociolek. Elsevier Inc. DOI: [10.1016/B978-0-12-385876-4.00015-3](https://doi.org/10.1016/B978-0-12-385876-4.00015-3).
- Komárek, J. and Johansen, J. R. (2015a). "Coccoid Cyanobacteria". In: *Freshwater algae of North America*. Ed. by J. D. Wehr, R. G. Sheath, and J. P. Kociolek. Elsevier Inc. DOI: [10.1016/B978-0-12-385876-4.00003-7](https://doi.org/10.1016/B978-0-12-385876-4.00003-7).

- Komárek, J. and Johansen, J. R. (2015b). "Filamentous Cyanobacteria". In: *Freshwater algae of North America*. Ed. by J. D. Wehr, R. G. Sheath, and J. P. Kociolek. Elsevier Inc. DOI: [10.1016/B978-0-12-385876-4.00004-9](https://doi.org/10.1016/B978-0-12-385876-4.00004-9).
- Ladd, C. and Lee, H. C. (2005). "The use of biological and botanical evidence in criminal investigations". In: *Forensic botany: principles and applications to criminal casework*. Ed. by H. Miller Coyle. CRC Press, pp. 97–115.
- Lee, R. E. (2018). *Phycology*. Cambridge university press.
- Levin, E. A., Morgan, R. M., Scott, K. R., and Jones, V. J. (2017). "The transfer of diatoms from freshwater to footwear materials: An experimental study assessing transfer, persistence, and extraction methods for forensic reconstruction". *Science & Justice* **57**(5), 349–360. DOI: [10.1016/j.scijus.2017.0](https://doi.org/10.1016/j.scijus.2017.0).
- Liu, M., Zhao, Y., Sun, Y., Li, Y., Wu, P., Zhou, S., and Ren, L. (2020). "Comparative study on diatom morphology and molecular identification in drowning cases". *Forensic science international* **317**, 110552. DOI: [10.1016/j.forsciint.2020.110552](https://doi.org/10.1016/j.forsciint.2020.110552).
- Lombardi, G. (1999). "The contribution of forensic geology and other trace evidence analysis to the investigation of the killing of Italian Prime Minister Aldo Moro". *Journal of forensic sciences* **44**(3), 634–642. DOI: [10.1520/JFS14523J](https://doi.org/10.1520/JFS14523J).
- Magni, P. A., Mohan, M., Vadiveloo, A., and Moheimani, N. R. (2020). "Transferability of Australian diatoms to clothing: Assessment of several extraction methods on different fabric types under laboratory conditions". *Forensic Science International* **312**, 110297. DOI: [10.1016/j.forsciint.2020.110297](https://doi.org/10.1016/j.forsciint.2020.110297).
- Magrey, A. H. and Raj, M. (2014). "Role of diatoms in forensic diagnosis of drowning cases from Jammu & Kashmir, India". *Biosci Biotech Res Comm* **7**(1), 72–7. DOI: [10.4172/2157-7145.1000181](https://doi.org/10.4172/2157-7145.1000181).
- Marra, A. C. (2023). "Taphonomical investigation applied to clandestine graves". *Atti della Accademia Peloritana dei Pericolanti. Classe di Scienze Fisiche, Matematiche e Naturali* **101**(S1), A6. DOI: [10.1478/AAPP.101S1A6](https://doi.org/10.1478/AAPP.101S1A6).
- Marra, A. C., Di Silvestro, G., and Somma, R. (2023). "Palaeontology applied to criminal investigation". *Atti della Accademia Peloritana dei Pericolanti. Classe di Scienze Fisiche, Matematiche e Naturali* **101**(S1), A4. DOI: [10.1478/AAPP.101S1A4](https://doi.org/10.1478/AAPP.101S1A4).
- Morabito, M., Mondello, F., and Somma, R. (2023). "Macrobotanic data implementing Forensic Geology investigations". *Atti della Accademia Peloritana dei Pericolanti. Classe di Scienze Fisiche, Matematiche e Naturali* **101**(S1), A13. DOI: [10.1478/AAPP.101S1A13](https://doi.org/10.1478/AAPP.101S1A13).
- Palmer, C. M. (1959). *Algae in water supplies: an illustrated manual on the identification, significance, and control of algae in water supplies*. Vol. 2. US Department of Health.
- Picozzi, M. and Intini, A. (2009). *Scienze forensi: teoria e prassi dell'investigazione scientifica*. UTET giuridica.
- Pollanen, M. S. (1998). *Forensic diatomology and drowning*. Elsevier Health Sciences.
- Pollanen, M. S., Cheung, C., and Chiasson, D. A. (1997). "The diagnostic value of the diatom test for drowning, I. Utility: a retrospective analysis of 771 cases of drowning in Ontario, Canada". *Journal of forensic sciences* **42**(2), 281–285. DOI: [10.1520/JFS14111J](https://doi.org/10.1520/JFS14111J).
- Prescott, G. W., Croasdale, H. T., and Vinyard, W. C. (1964). "How to know the freshwater algae". *How to know the freshwater algae*.
- Prescott, G. W., Croasdale, H. T., Vinyard, W. C., and M. Bicudo, C. E. de (1975). *A Synopsis of North American Desmids: Part II: Desmidiaceae: Placodermae. Section 3*. University of Nebraska Press.
- Ruffell, A. and McKinley, J. (2009). *Geoforensics*. John Wiley & Sons Ltd.
- Saferstein, R. (2017). *Criminalistics: An introduction to forensic science*.

- Saini, E., Khanagwal, V. P., and Singh, R. (2017). "A systematic databasing of diatoms from different geographical localities and sites of Haryana for advancing validation of forensic diatomology". *Data in brief* **10**, 63–68. DOI: [10.1016/j.dib.2016.11.072](https://doi.org/10.1016/j.dib.2016.11.072).
- Scott, K. R., Jones, V. J., Cameron, N. G., Young, J. M., and Morgan, R. M. (2021). "Freshwater diatom persistence on clothing I: A quantitative assessment of trace evidence dynamics over time". *Forensic Science International* **325**, 110898. DOI: [10.1016/j.forsciint.2021.110898](https://doi.org/10.1016/j.forsciint.2021.110898).
- Scott, K. R., Morgan, R. M., Jones, V. J., and Cameron, N. G. (2014). "The transferability of diatoms to clothing and the methods appropriate for their collection and analysis in forensic geoscience". *Forensic science international* **241**, 127–137. DOI: [10.1016/j.forsciint.2014.05.011](https://doi.org/10.1016/j.forsciint.2014.05.011).
- Scott, K. R., Morgan, R. M., Jones, V. J., Dudley, A., Cameron, N., and Bull, P. A. (2017). "The value of an empirical approach for the assessment of diatoms as environmental trace evidence in forensic limnology". *Archaeological and Environmental Forensic Science* **1**(1), 49–78. DOI: [10.1558/aefs.32474](https://doi.org/10.1558/aefs.32474).
- Sharma, M. K., Sharma, A., and Bhatt, A. K. (2021). "Diatom, a phytoplanktonic community study from different water bodies of District Hamirpur, Himachal Pradesh: Role in forensic studies and biotechnological innovations". *International Journal of Lakes and Rivers* **14**(1), 93–111.
- Sheath, R. G. and Wehr, J. D. (2015). "Introduction to the freshwater algae". In: *Freshwater Algae of North America*. Ed. by J. D. Wehr, R. G. Sheath, and J. P. Kociolek, pp. 1–11. DOI: [10.1016/B978-0-12-385876-4.00001-3](https://doi.org/10.1016/B978-0-12-385876-4.00001-3).
- Shen, X., Liu, Y., Xiao, C., Zheng, C., Huang, J., Shi, H., Xu, Q., Cheng, J., Liu, C., and Zhao, J. (2019). "Analysis of false-positive results of diatom test in the diagnosis of drowning—would not be an impediment". *International journal of legal medicine* **133**, 1819–1824. DOI: [10.1007/s00414-019-02021-4](https://doi.org/10.1007/s00414-019-02021-4).
- Shubert, E. and Gärtner, G. (2015). "Nonmotile coccoid and colonial green algae". In: *Freshwater Algae of North America*. Ed. by J. D. Wehr, R. G. Sheath, and J. P. Kociolek. Elsevier, pp. 315–373. DOI: [10.1016/B978-0-12-385876-4.00007-4](https://doi.org/10.1016/B978-0-12-385876-4.00007-4).
- Siver, P. A., Lord, W. D., and McCarthy, D. J. (1994). "Forensic limnology: the use of freshwater algal community ecology to link suspects to an aquatic crime scene in southern New England". *Journal of Forensic Sciences* **39**, 847–847. DOI: [10.1016/B978-0-12-385876-4.00007-4](https://doi.org/10.1016/B978-0-12-385876-4.00007-4).
- Somma, R. (2022). "Advances in Flipped Classrooms for Teaching and Learning Forensic Geology". *Education Sciences* **12**(6), 403. DOI: [10.3390/educsci12060403](https://doi.org/10.3390/educsci12060403).
- Somma, R. (2023a). "A multidisciplinary approach based on the cooperation of forensic geologists, botanists, and engineers: Computed Axial Tomography applied to a case work". *Atti della Accademia Peloritana dei Pericolanti. Classe di Scienze Fisiche, Matematiche e Naturali* **101**(S1), A12. DOI: [10.1478/AAPP.101S1A12](https://doi.org/10.1478/AAPP.101S1A12).
- Somma, R. (2023b). "The space and time dimensions in the criminal behaviour of lust murderers". *Atti della Accademia Peloritana dei Pericolanti. Classe di Scienze Fisiche, Matematiche e Naturali* **101**(S1), A19. DOI: [10.1478/AAPP.101S1A19](https://doi.org/10.1478/AAPP.101S1A19).
- Somma, R. (2023c). "Unraveling crimes with geosciences". *Atti della Accademia Peloritana dei Pericolanti. Classe di Scienze Fisiche, Matematiche e Naturali* **101**(S1), A2. DOI: [10.1478/AAPP.101S1A2](https://doi.org/10.1478/AAPP.101S1A2).
- Somma, R., Altadonna, A., Cucinotta, F., Raffaele, M., Salmeri, F., Baldino, G., Ventura Spagnolo, E., and Sapienza, D. (2023a). "The technologies of Laser Scanning and Structured Blue Light Scanning applied to criminal investigation: case studies". *Atti della Accademia Peloritana dei Pericolanti. Classe di Scienze Fisiche, Matematiche e Naturali* **101**(S1), A15. DOI: [10.1478/AAPP.101S1A15](https://doi.org/10.1478/AAPP.101S1A15).
- Somma, R., Baldino, G., Altadonna, A., Asmundo, A., Fodale, V., Gualniera P. abd Mondello, C., Pennisi, C., Raffaele, M., Salmeri, F., Ventura Spagnolo, E., and Sapienza, D. (2023b). "Education and training activities in forensic and biomedical sciences: The Laser scanner technology". *Atti*

- della Accademia Peloritana dei Pericolanti. Classe di Scienze Fisiche, Matematiche e Naturali* **101**(S1), A16. DOI: [10.1478/AAPP.101S1A16](https://doi.org/10.1478/AAPP.101S1A16).
- Somma, R., Cascio, M., Cucinotta, F., Mondello, F., and Morabito, M. (2023c). “Recent advances in forensic geology and botany for the reconstruction of event dynamics in outdoor crime scenes: a case study”. *Atti della Accademia Peloritana dei Pericolanti. Classe di Scienze Fisiche, Matematiche e Naturali* **101**(S1), A10. DOI: [10.1478/AAPP.101S1A10](https://doi.org/10.1478/AAPP.101S1A10).
- Somma, R. and Costa, N. (2022). “Unraveling Crimes with Geology: As Geological and Geographical Evidence Related to Clandestine Graves May Assist the Judicial System”. *Geosciences* **12**(9), 339. DOI: [10.3390/geosciences12090339](https://doi.org/10.3390/geosciences12090339).
- Somma, R. and Costa, N. (2023). “GIS-based RAG-coded search priority scenarios for predictive maps to prevent future serial serious crimes: the case study of the Florence Monster”. *Atti della Accademia Peloritana dei Pericolanti. Classe di Scienze Fisiche, Matematiche e Naturali* **101**(S1), A18. DOI: [10.1478/AAPP.101S1A18](https://doi.org/10.1478/AAPP.101S1A18).
- Somma, R. and Maniscalco, R. (2023). “Forensic geology applied to criminal investigation: a case report”. *Atti della Accademia Peloritana dei Pericolanti. Classe di Scienze Fisiche, Matematiche e Naturali* **101**(S1), A9. DOI: [10.1478/AAPP.101S1A9](https://doi.org/10.1478/AAPP.101S1A9).
- Somma, R., Spoto, S. E., Raffaele, M., and Salmeri, F. (2023d). “Measuring color techniques for forensic comparative analyses of geological evidence”. *Atti della Accademia Peloritana dei Pericolanti. Classe di Scienze Fisiche, Matematiche e Naturali* **101**(S1), A14. DOI: [10.1478/AAPP.101S1A14](https://doi.org/10.1478/AAPP.101S1A14).
- Somma, R., Sutton, L., and Byrd, J. H. (2023). “Forensic geology applied to the search for homicide graves”. *Atti della Accademia Peloritana dei Pericolanti. Classe di Scienze Fisiche, Matematiche e Naturali* **101**(S1), A5. DOI: [10.1478/AAPP.101S1A5](https://doi.org/10.1478/AAPP.101S1A5).
- Spikmans, V. (2019). “The evolution of environmental forensics: From laboratory to field analysis”. *WIREs Forensic Science* **1**(3), e1334. DOI: <https://doi.org/10.1002/wfs2.1334>.
- Spoto, S. E. (2023). “Illicit trafficking of diamonds: new frontiers”. *Atti della Accademia Peloritana dei Pericolanti. Classe di Scienze Fisiche, Matematiche e Naturali* **101**(S1), A3. DOI: [10.1478/AAPP.101S1A3](https://doi.org/10.1478/AAPP.101S1A3).
- Spoto, S. E., Somma, R., and Crea, F. (2021). “Using a forensic-based learning approach to teach geochemistry”. *Atti della Accademia Peloritana dei Pericolanti. Classe di Scienze Fisiche, Matematiche e Naturali* **99**(S1), A31. DOI: [10.1478/AAPP.99S1A31](https://doi.org/10.1478/AAPP.99S1A31).
- Spoto, S. E., Barone, S., and Somma, R. (2023). “An introduction to forensic geosciences”. *Atti della Accademia Peloritana dei Pericolanti. Classe di Scienze Fisiche, Matematiche e Naturali* **101**(S1), A1. DOI: [10.1478/AAPP.101S1A1](https://doi.org/10.1478/AAPP.101S1A1).
- Stevenson, R. J., Bothwell, M. L., Lowe, R. L., and Thorp, J. H. (1996). *Algal ecology: Freshwater benthic ecosystem*. Academic press. DOI: [10.1016/B978-0-12-668450-6.X5027-9](https://doi.org/10.1016/B978-0-12-668450-6.X5027-9).
- Tagliabue, G., Masseroli, A., Mattia, M., Sala, C., Belgiovine, E., Capuzzo, D., Galimberti, P., Slavazzi, F., Cattaneo, C., and Trombino, L. (2023). “Thanatogenic Anthrosols: a geoforensic approach to the exploration of the Sepolcreto of the Ca’ Granda (Milan)”. *Atti della Accademia Peloritana dei Pericolanti. Classe di Scienze Fisiche, Matematiche e Naturali* **101**(S1), A8. DOI: [10.1478/AAPP.101S1A8](https://doi.org/10.1478/AAPP.101S1A8).
- Thakar, M., Luthra, D., and Khattar, J. I. S. (2018). “Forensic studies of phytoplankton ecology of two water bodies of Kurukshetra area of Haryana, State in India”. *Egyptian Journal of Forensic Sciences* **8**, 0068-4. DOI: [10.1186/s41935-018-0068-4](https://doi.org/10.1186/s41935-018-0068-4).
- Trainor, F. R. (2014). *Biological Aspects of Scenedesmus (Chlorophyceae)—Phenotypic Plasticity*. Nova Hedwigia, J. Cramer.
- Verbruggen, H. and Theriot, E. C. (2008). “Building trees of algae: some advances in phylogenetic and evolutionary analysis”. *European Journal of Phycology* **43**(3), 229–252. DOI: [10.1080/09670260802207530](https://doi.org/10.1080/09670260802207530).

- Wanner, M., Betker, E., Shimano, S., and Krawczynski, R. (2018). “Are soil testate amoebae and diatoms useful for forensics?” *Forensic Science International* **289**, 223–231. DOI: [10.1016/j.forsciint.2018.05.027](https://doi.org/10.1016/j.forsciint.2018.05.027).
- Wehr, J. D., Sheath, R. G., and Kociolek, J. P. (2015). *Freshwater Algae of North America: Ecology and Classification*. Elsevier. DOI: [10.1016/C2010-0-66664-8](https://doi.org/10.1016/C2010-0-66664-8).
- Zhang, P., Kang, X., Zhang, S., Xiao, C., Ma, Y., Shi, H., Xu, Q., Zhao, J., Chen, L., and Liu, C. (2020). “The length and width of diatoms in drowning cases as the evidence of diatoms penetrating the alveoli-capillary barrier”. *International Journal of Legal Medicine* **134**, 1–6. DOI: [10.1007/s00414-019-02164-4](https://doi.org/10.1007/s00414-019-02164-4).
- Zhao, J., Ma, Y., Liu, C., Wen, J., Hu, S., Shi, H., and Zhu, L. (2016). “A quantitative comparison analysis of diatoms in the lung tissues and the drowning medium as an indicator of drowning”. *Journal of Forensic and Legal Medicine* **42**, 75–78. DOI: [10.1016/j.jflm.2016.05.021](https://doi.org/10.1016/j.jflm.2016.05.021).
- Zhou, Y., Cao, Y., Huang, J., Deng, K., Ma, K., Zhang, T., Chen, L., Zhang, J., and Huang, P. (2020). “Research advances in forensic diatom testing”. *Forensic Sciences Research* **5**, 1–8. DOI: [10.1080/20961790.2020.1718901](https://doi.org/10.1080/20961790.2020.1718901).
- Zimmerman, K. and Wallace, J. R. (2008). “The Potential to Determine a Postmortem Submersion Interval Based on Algal/Diatom Diversity on Decomposing Mammalian Carcasses in Brackish Ponds in Delaware”. *Journal of forensic sciences* **53**, 935–941. DOI: [10.1111/j.1556-4029.2008.00748.x](https://doi.org/10.1111/j.1556-4029.2008.00748.x).

^a Università degli Studi di Messina,
Dipartimento di Scienze Chimiche, Biologiche, Farmaceutiche e Ambientali,
Viale F. Stagno d’Alcontres 31, 98166 Messina, Italy

^b Accademia Peloritana dei Pericolanti,
Palazzo Università,
Piazza Pugliatti 1, 98122 Messina, Italy

^c Università degli Studi di Messina,
Dipartimento di Scienze Matematiche e Informatiche, Scienze Fisiche e Scienze della Terra,
Viale F. Stagno d’Alcontres 31, 98166 Messina, Italy

* To whom correspondence should be addressed | email: marina.morabito@unime.it

Paper contributed to the workshop on “Advances and applications in geoforensics: Unraveling crimes with geology”,
held in Messina, Italy (26 September 2022) under the patronage of the *Accademia Peloritana dei Pericolanti*

Manuscript received 17 March 2023; published online 12 September 2023



© 2023 by the author(s); licensee *Accademia Peloritana dei Pericolanti* (Messina, Italy). This article is an open access article distributed under the terms and conditions of the [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) (<https://creativecommons.org/licenses/by/4.0/>).